

A Note on the Geometry of Green-Lazarsfeld Sets

Alexander Brudnyi*

Department of Mathematics and Statistics
University of Calgary
Calgary, Canada

Abstract

We study some subsets of the Green-Lazarsfeld set $\Sigma^1(M)$ for a compact Kähler manifold M .

1. Introduction.

Let M be a compact Kähler manifold and $H^1(M, \mathbb{C}^*) := \text{Hom}(\pi_1(M), \mathbb{C}^*)$. Given a character $\xi \in H^1(M, \mathbb{C}^*)$ let $\mathbb{C}_\xi$ denote the associated $\pi_1(M)$ -module. Let $\Sigma^1(M)$ be the set of characters ξ such that $H^1(\pi_1(M), \mathbb{C}_\xi)$ is nonzero. The structure of $\Sigma^1(M)$ was described in the papers of Beauville [Be], Simpson [S], and Campana [Ca]:

There is a finite number of surjective holomorphic maps with connected fibres $f_i : M \rightarrow C_i$ onto smooth compact complex curves of genus $g \geq 1$ and torsion characters $\rho_i, \xi_j \in H^1(M, \mathbb{C}^)$ such that*

$$\Sigma^1(M) = \bigcup_i \rho_i f_i^* H^1(C_i, \mathbb{C}^*) \cup \bigcup_j \{\xi_j\}.$$

In this paper we study the components $f_i^* H^1(C_i, \mathbb{C}^*)$ corresponding to trivial ρ_i .

Theorem 1.1 *Let $f : M \rightarrow C$ be a surjective holomorphic map with connected fibres of a compact Kähler manifold M onto a smooth complex curve C of genus $g \geq 1$. Assume that $\xi \in H^1(C, \mathbb{C}^*)$ is not torsion. Then*

$$f^* : H^1(\pi_1(C), \mathbb{C}_\xi) \longrightarrow H^1(\pi_1(M), \mathbb{C}_{f^*\xi})$$

is an isomorphism.

*1991 *Mathematics Subject Classification*. Primary 20F34. Secondary 58A10.
Key words and phrases. Kähler manifold, cohomology group, torsion character.

2. Proof of Theorem 1.1.

2.1. Let $T_2(\mathbb{C}) \subset GL_2(\mathbb{C})$ be the Lie subgroup of upper triangular matrices. By $T_2 \subset T_2(\mathbb{C})$ we denote the Lie group of matrices of the form

$$\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \quad (a \in \mathbb{C}^*, b \in \mathbb{C}).$$

Also by $D_2 \subset T_2$ we denote the subgroup of diagonal matrices and by $N_2 \subset T_2$ the subgroup of unipotent matrices. Let M be a compact Kähler manifold. For a homomorphism $\rho \in \text{Hom}(\pi_1(M), T_2)$ we let $\rho_a \in \text{Hom}(\pi_1(M), \mathbb{C}^*)$ denote the upper diagonal character of ρ . Further, N_2 acts on T_2 by conjugation. Any two homomorphisms from $\text{Hom}(\pi_1(M), T_2)$ belonging to the orbit of this action will be called *equivalent*. It is well known that the class of equivalence of ρ is uniquely defined by an element $c_\rho \in H^1(\pi_1(M), \mathbb{C}_{\rho_a})$ (see e.g. [A, Prop. 2]). Now our theorem is a consequence of the following result.

Theorem 2.1 *Assume that $\rho \in \text{Hom}(\pi_1(M), T_2)$ satisfies $\text{Ker}(f_*) \subset \text{Ker}(\rho_a)$ but $\text{Ker}(f_*) \not\subset \text{Ker}(\rho)$. Then ρ_a is a torsion character.*

The next several sections contain some results used in the proof of Theorem 2.1.

2.2. Let $f : M \rightarrow C$ be a holomorphic surjective map with connected fibres of a compact Kähler manifold M onto a smooth compact complex curve C . Consider a flat vector bundle L on C of complex rank 1 with unitary structure group and let $E = f^*L$ be the pullback of this bundle to M . Let $\omega \in \Omega^1(E)$ be a holomorphic 1-form with values in E (by the Hodge decomposition ω is d -closed).

Lemma 2.2 *Assume that $\omega|_{V_z} = 0$ for the fibre $V_z := f^{-1}(z)$ over a regular value $z \in C$. Then $\omega|_V = 0$ for any fibre V of f .*

If V is not smooth the lemma asserts that ω equals 0 on the smooth part of V .

Proof. Denote by $S \subset C$ the finite set of non-regular values of f . By Sard's theorem, $f : M \setminus f^{-1}(S) \rightarrow C \setminus S$ is a fibre bundle with connected fibres. According to our assumption there is a fibre V_z of this bundle such that $\omega|_{V_z} = 0$. Then $\omega|_V = 0$ for any fibre V of $f|_{M \setminus f^{-1}(S)}$. In fact, for any fibre V there is an open neighbourhood O_V of V diffeomorphic to $\mathbb{R}^2 \times V$ such that $E|_{O_V} (\cong E|_V)$ is the trivial flat vector bundle (here $E|_V$ is trivial as the pullback of a bundle defined over a point). Any d -closed holomorphic 1-form defined on O_V that vanishes on V is d -exact, and so it vanishes on each fibre contained in O_V . Starting with a tubular neighbourhood O_{V_z} and taking into account that $\omega|_{O_{V_z}}$ can be considered as a d -closed holomorphic 1-form (because $E|_{O_{V_z}}$ is trivial) we obtain that $\omega|_V = 0$ for any fibre $V \subset O_{V_z}$. Then the required statement follows by induction if we cover $M \setminus f^{-1}(S)$ by open tubular neighbourhoods of fibres of f and use the fact that $C \setminus S$ is connected.

Consider now fibres over the singular part S . Let $O_x \subset C$ be the neighbourhood of a point $x \in S$ such that $O_x \setminus S = O_x \setminus \{x\}$ is biholomorphic to $\mathbb{D} \setminus \{0\}$. In particular, $\pi_1(O_x \setminus S) = \mathbb{Z}$. Without loss of generality we may assume that $V = f^{-1}(x)$ is a deformation retract of an open neighbourhood U_x of V and $W_x := f^{-1}(O_x) \subset U_x$

(such U_x exists, e.g., by the triangulation theorem of the pair (M, V) , see [L, Th.2]). Since $E|_V$ is trivial, the bundle $E|_{U_x}$ is also trivial. Therefore we can regard ω as a d -closed holomorphic 1-form on U_x . Moreover, we already have proved that $\omega|_F = 0$ for any fibre $F \subset W_x \setminus f^{-1}(S)$. So ω equals 0 on each fibre of the fibration $f : W_x \setminus f^{-1}(S) \rightarrow O_x \setminus S$. This implies that there is a d -closed holomorphic 1-form ω_1 on $O_x \setminus S$ such that $\omega = f^*(\omega_1)$. Assume now that $\omega|_V \neq 0$. Then integration of $\omega|_{W_x}$ along paths determines a non-trivial homomorphism $h_{\omega,x} : \pi_1(W_x) \rightarrow \mathbb{C}$ whose image is isomorphic to $\mathbb{Z}$. Indeed, since embedding $W_x \setminus f^{-1}(S) \hookrightarrow W_x$ induces a surjective homomorphism of fundamental groups, we can integrate ω by paths contained in $W_x \setminus f^{-1}(S)$ obtaining the same image in $\mathbb{C}$. Further, for any path $\gamma \subset W_x \setminus f^{-1}(S)$ we have $\int_\gamma \omega = \int_{\gamma_1} \omega_1$, where γ_1 is a path in $O_x \setminus S$ representing the element $f_*(\gamma) \in \pi_1(O_x \setminus S)$. But $\pi_1(O_x \setminus S) = \mathbb{Z}$ and therefore $h_{\omega,x}(\pi_1(W_x)) \cong \mathbb{Z}$. Now, any path in W_x is homotopically equivalent inside U_x to a path contained in V and so we can define $h_{\omega,x}$ integrating $\omega|_V$. Thus we obtain a homomorphism h of $\Gamma := (\pi_1(V)/D\pi_1(V))/_{torsion}$ into $\mathbb{C}$ with the same image as for $h_{\omega,x}$, i.e., $\cong \mathbb{Z}$. Without loss of generality we may assume that V is smooth (for otherwise, we apply the arguments below to each irreducible component of a desingularization of V). Integrating holomorphic 1-forms on V along paths we embed Γ into some $\mathbb{C}^k$ as a lattice of rank $2k$. Then there is a linear holomorphic functional f_ω on $\mathbb{C}^k$ such that $h = f_\omega|_\Gamma$. Since $rk(h(\Gamma)) = 1$, f_ω equals 0 on a subgroup $H \subset \Gamma$ isomorphic to $\mathbb{Z}^{2k-1}$. In particular, $f_\omega = 0$ on the vector space $H \otimes \mathbb{R}$ of real dimension $2k-1$. But $Ker(f_\omega)$ is a complex vector space. So $f_\omega = 0$ on $\mathbb{C}^k$ which implies that $\omega|_V = 0$. This contradicts our assumption and proves the required statement for fibres over the points of S .

The lemma is proved. $\square$

2.3. Next we prove that the character ρ_a in Theorem 2.1 cannot be non-unitary.

Proposition 2.3 *Let $\rho \in Hom(\pi_1(M), T_2)$ be such that ρ_a is a non-unitary character. Assume that $\rho_a|_{Ker(f_*)}$ is trivial. Then $\rho|_{Ker(f_*)}$ is also trivial.*

Proof. Let $p_t : M_t \rightarrow M$ be the Galois covering with transformation group $Tor(\pi_1(M)/D\pi_1(M))$. Then $\rho|_{\pi_1(M_t)}$ determines a C^∞ -trivial complex rank-2 flat vector bundle on M_t , because $\rho_a|_{\pi_1(M_t)} = \exp(\rho'|_{\pi_1(M_t)})$ for some $\rho' \in Hom(\pi_1(M), \mathbb{C})$. In particular, $\rho|_{\pi_1(M_t)}$ can be defined by a flat connection

$$\Omega = \begin{pmatrix} \omega & \eta \\ 0 & 0 \end{pmatrix}; \quad d\Omega - \Omega \wedge \Omega = 0$$

on $M_t \times \mathbb{C}^2$ where ω is a d -harmonic 1-form on M_t lifted from M . Moreover, since $\rho_a|_{Ker(f_*)}$ is trivial, ω is the pullback by $f \circ p_t$ of a d -harmonic 1-form defined on C . Let $\omega = \omega_1 + \omega_2$ be the type decomposition of ω into the sum of holomorphic and antiholomorphic 1-forms lifted from C . Denote by E_ρ the rank-1 flat vector bundle on M_t with unitary structure group constructed by the flat connection $\omega_2 - \overline{\omega_2}$ and by $E_0 = M_t \times \mathbb{C}$ the trivial flat vector bundle. Observe that E_ρ is pullback of a flat bundle on C . In particular, $E_\rho|_V$ is trivial for any fibre V of $f \circ p_t$. According to [Br, Th.1.2] the equivalence class of $\rho|_{\pi_1(M_t)}$ is determined by a d -harmonic 1-form θ

with values in $\text{End}(E_\rho \oplus E_0)$ satisfying $\theta \wedge \theta$ represents 0 in $H^2(M_t, \text{End}(E_\rho \oplus E_0))$. More precisely,

$$\theta = \begin{pmatrix} \omega_1 & \eta' \\ 0 & 0 \end{pmatrix}$$

where η' is a d -harmonic 1-form with values in E_ρ satisfying $\omega_1 \wedge \eta'$ represents 0 in $H^2(M_t, E_\rho)$. Let $M_t \xrightarrow{g_1} M_1 \xrightarrow{g_2} C$ be the Stein factorization of $f \circ p_t$. Here g_1 is a morphism with connected fibres onto a smooth curve M_1 and g_2 is a finite morphism. Then for any fibre $V \hookrightarrow M_t$ of g_1 we have that $p_t|_V : V \rightarrow p_t(V)$ is a regular covering of the fibre $p_t(V)$ of f with a finite abelian transformation group.

Lemma 2.4 $\eta'|_V = 0$ for fibres over regular values of g_1 .

Based on this statement we, first, finish the proof of the proposition and then will prove the lemma. From the lemma and Lemma 2.2 it follows that $\eta'|_W = 0$ for any fibre $i : W \hookrightarrow M_t$ of g_1 . Thus $\rho|_{\pi_1(M_t)}$ is trivial on $i_*(\pi_1(W)) \subset \pi_1(M_t)$. But $\pi_1(W)$ is a subgroup of a finite index in $\pi_1(p_t(W))$ and the image of $\rho|_{j_*(\pi_1(p_t(W)))}$ consists of unipotent matrices by the assumption of the proposition. (Here $j : p_t(W) \hookrightarrow M$ is embedding.) Therefore $\rho|_{j_*(\pi_1(p_t(W)))}$ is trivial for any W . Denote by E the flat vector bundle on M associated to ρ . Then we have proved that $E|_W$ is trivial for any fibre W of f . Further, let $(U_i)_{i \in I}$ be an open cover of C such that $W_i := f^{-1}(U_i)$ is an open neighbourhood of a fibre $V_i = f^{-1}(x_i)$, $x_i \in U_i$, and W_i is deformable onto V_i . Since $E|_{V_i}$ is trivial, $E|_{W_i}$ is also trivial. In particular, E is defined by a locally constant cocycle $\{c_{ij}\}$ defined on the cover $(W_i)_{i \in I}$. But then $\{c_{ij}\}$ is the pullback of a cocycle defined on $(U_i)_{i \in I}$ because the fibres of f are connected. This cocycle determines a bundle E' on C such that $f^*E' = E$. Then ρ is the pullback of a homomorphism $\rho' \in \text{Hom}(\pi_1(C), T_2)$ constructed by E' .

This completes the proof of the proposition modulo Lemma 2.4.

Proof of Lemma 2.4. Observe that $\omega_1 \neq 0$ in the above definition of θ because ρ_a is non-unitary. We also regard $\eta' \neq 0$, for otherwise, the image of ρ consists of diagonal matrices and the required statement is trivial. Let $i : V \hookrightarrow M_t$ be a fibre over a regular value of g_1 . For any $\lambda \in \mathbb{C}^*$ consider the form

$$\theta_\lambda = \begin{pmatrix} \lambda\omega_1 & \eta' \\ 0 & 0 \end{pmatrix}.$$

Clearly, $\theta_\lambda \wedge \theta_\lambda$ represents 0 in $H^2(M_t, \text{End}(E_\rho \oplus E_0))$ and thus, by [Br, Prop.2.4], it determines a representation $\rho_\lambda \in \text{Hom}(\pi_1(M_t), T_2)$ with the upper diagonal character $\rho_{\lambda a}$ defined by the flat connection $\lambda\omega_1 + \omega_2$. In particular, the family $\{\rho_{\lambda a}\}$ contains infinitely many different characters. Assume that $\rho|_{i_*(\pi_1(V))}$ is not trivial. Then it can be determined by the restriction $\theta|_{i(V)}$. But according to our assumption $\omega_1|_{i(V)} = 0$ and $E_\rho|_{i(V)}$ is a trivial flat vector bundle. Thus $\rho|_{i_*(\pi_1(V))}$ is defined by $\eta'|_{i(V)} \in H^1(i(V), \mathbb{C})$. Let $\psi \in \text{Hom}(i(V), \mathbb{C})$ be a homomorphism obtained by integration of $\eta'|_{i(V)}$ along paths generating $\pi_1(i(V))$. Then

$$\rho|_{i_*(\pi_1(V))} = \begin{pmatrix} 1 & \psi \\ 0 & 1 \end{pmatrix}.$$

We also obtain that $\rho_\lambda|_{i_*(\pi_1(V))} = \rho|_{i_*(\pi_1(V))}$ and so it is not trivial for any $\lambda \in \mathbb{C}^*$. We will prove now that the family $\{\rho_{\lambda a}\}$ contains finitely many different characters. This contradicts our assumption and shows that $\rho|_{i_*(\pi_1(V))}$ is trivial and so $\eta'|_V = 0$.

Let $H := i_*(\pi_1(V))$ be a normal subgroup of $\pi_1(M_t)$ (the normality follows from the exact homotopy sequence for the bundle defined over regular values of g_1). Denote by H^{ab} the quotient H/DH . Then H^{ab} is an abelian group of a finite rank. Moreover, H^{ab} is a normal subgroup of $K := \pi_1(M_t)/DH$ and the group $K_1 := K/H^{ab}$ is finitely generated. Note that ρ_λ induces a homomorphism $\hat{\rho}_\lambda : K \rightarrow T_2$ for any $\lambda \in \mathbb{C}^*$ with the same image as for ρ_λ because the image of $\rho_\lambda|_H$ is abelian. In particular, $\hat{\rho}_\lambda(H^{ab})$ is a normal subgroup of $\hat{\rho}(K)$. Since by our assumption $\hat{\rho}_\lambda(H^{ab})$ is a non-trivial subgroup of unipotent matrices, from the identity

$$\begin{pmatrix} c & b \\ 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} 1 & v \\ 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} c & b \\ 0 & 1 \end{pmatrix}^{-1} = \begin{pmatrix} 1 & c \cdot v \\ 0 & 1 \end{pmatrix} \quad (c \in \mathbb{C}^*, b, v \in \mathbb{C}) \quad (2.1)$$

it follows that the action of $\hat{\rho}_\lambda(K)$ on $\hat{\rho}_\lambda(H^{ab})$ by conjugation is defined by multiplication of non-diagonal elements of $\hat{\rho}_\lambda(H^{ab})$ by elements of $\rho_{\lambda a}(\pi_1(M_t))$. Note also that $Tor(H^{ab})$ belongs to $Ker(\hat{\rho}_\lambda)$ for any λ . Thus if $H' \cong \mathbb{Z}^s$ is a maximal free abelian subgroup of H^{ab} then $\rho_{\lambda a}(\pi_1(M_t))$ consists of eigen values of matrices from $SL_s(\mathbb{Z})$ obtained by the natural action (by conjugation) of K_1 on H' . Since K_1 is finitely generated, the number of different $\rho_{\lambda a}$ is finite which is false.

This contradiction completes the proof of the lemma. $\square$

2.4. Let $f : M \rightarrow C$ be a holomorphic map with connected fibres of a compact Kähler manifold M onto a smooth compact complex curve C of genus $g \geq 1$. Then $\pi_1(M)$ is defined by the exact sequence

$$\{1\} \longrightarrow Ker(f_*) \longrightarrow \pi_1(M) \longrightarrow \pi_1(C) \longrightarrow \{1\}.$$

Let $G \subset Ker(f_*)$ be a normal subgroup of $\pi_1(M)$. The quotient $R := \pi_1(M)/G$ is defined by the sequence

$$\{1\} \longrightarrow Ker(f_*)/G \longrightarrow R \longrightarrow \pi_1(C) \longrightarrow \{1\}.$$

Assume that

- (1) $H := Ker(f_*)/G$ is a free abelian group of finite rank $k \geq 1$;
- (2) the natural action of $D\pi_1(C)$ on H is trivial.

From (2) it follows that the action s of $\pi_1(C)$ on H determines an action of $\mathbb{Z}^{2g} \cong \pi_1(C)/D\pi_1(C)$ on H . Identifying H with $\mathbb{Z}^k$ we can think of H as a subgroup (lattice) of $\mathbb{Z}^k \otimes \mathbb{C} = \mathbb{C}^k$. Then s determines a representation $s' : \pi_1(C) \rightarrow GL_k(\mathbb{C})$ such that $D\pi_1(C) \subset Ker(s')$ and $s'(g)|_H = s(g)$ for any $g \in \pi_1(C)$. Since s' descends to a representation $\mathbb{Z}^{2g} \rightarrow GL_k(\mathbb{C})$, it admits a decomposition $s' = \bigoplus_{j=1}^m s_j$ where s_j is equivalent to a nilpotent representation $\pi_1(C) \rightarrow T_{k_i}(\mathbb{C})$ with a diagonal character ρ_j . Here $\sum_{j=1}^m k_j = k$.

Proposition 2.5 *All characters ρ_j are torsion.*

Proof. By definition, R is an extension of $\pi_1(C)$ by H . It is well known, see, e.g. [G, Ch.I, Sec.6], that the class of extensions equivalent to R is uniquely defined by an element $c \in H^2(\pi_1(C), H)$ where the cohomology is defined by the action s of $\pi_1(C)$ on H . Let $f \in Z^2(\pi_1(C), H)$ be a cocycle determining c . Then one can define a representative of the equivalence class of extensions as the direct product $H \times \pi_1(C)$ with multiplication

$$(h_1, g_1) \cdot (h_2, g_2) = (h_1 + s(g_1)(h_2) + f(g_1, g_2), g_1 \cdot g_2);$$

$$h_1, h_2 \in H, \quad g_1, g_2 \in \pi_1(C) .$$

The natural embedding $H \hookrightarrow \mathbb{Z}^k \otimes \mathbb{C} (= \mathbb{C}^k)$ determines an embedding i of R into the group R' defined as $\mathbb{C}^k \times \pi_1(C)$ with multiplication

$$(v_1, g_1) \cdot (v_2, g_2) = (v_1 + s'(g_1)(v_2) + f(g_1, g_2), g_1 \cdot g_2);$$

$$f \in Z^2(\pi_1(C), H), \quad v_1, v_2 \in \mathbb{C}^k, \quad g_1, g_2 \in \pi_1(C) .$$

Here we regard f as an element of $Z^2(\pi_1(C), \mathbb{C}^k)$ defined by the action s' . From the decomposition $s' = \bigoplus_{j=1}^m s_j$ it follows that there is an invariant $\pi_1(C)$ -submodule $V_j \subset \mathbb{C}^k$ of $\dim_{\mathbb{C}} V_j = k - 1$ such that $W_j = \mathbb{C}^k / V_j$ is a one-dimensional $\pi_1(C)$ -module and the action of $\pi_1(C)$ on W_j is defined as multiplication by the character ρ_j . Then, by definition, V_j is a normal subgroup of R' and the quotient group $R_j = R' / V_j$ is defined by the sequence

$$\{1\} \longrightarrow \mathbb{C} \longrightarrow R_j \longrightarrow \pi_1(C) \longrightarrow \{1\} .$$

Here the action of $\pi_1(C)$ on $\mathbb{C}$ is multiplication by the character ρ_j . Further, the equivalence class of extensions isomorphic to R_j is defined by an element $c_j \in H^2(\pi_1(C), \mathbb{C})$ (the cohomology is defined by the above action of $\pi_1(C)$ on $\mathbb{C}$). We will assume that the character ρ_j is non-trivial (for otherwise, ρ_j is clearly torsion). Let us denote by t_j the composite homomorphism $\pi_1(M) \longrightarrow R \xrightarrow{i} R' \longrightarrow R_j$.

Let E_{ρ_j} be a complex rank-1 flat vector bundle on C constructed by $\rho_j \in \text{Hom}(\pi_1(C), \mathbb{C}^*)$. Since C is a $K(\pi_1(C), 1)$ -space, there is a natural isomorphism of the above group $H^2(\pi_1(C), \mathbb{C})$ and the Čech cohomology group $H^2(C, \mathbf{E}_{\rho_j})$ of the sheaf of locally constant sections of E_{ρ_j} , see e.g. [M, Ch.1, Complement to Sec.2]. But each flat vector bundle on C is C^∞ -trivial and each homomorphism from $\text{Hom}(\pi_1(C), \mathbb{C}^*)$ can be continuously deformed inside of $\text{Hom}(\pi_1(C), \mathbb{C}^*)$ to the trivial homomorphism. Therefore by the index theorem

$$\dim_{\mathbb{C}} H^0(C, \mathbf{E}_{\rho_j}) - \dim_{\mathbb{C}} H^1(C, \mathbf{E}_{\rho_j}) + \dim_{\mathbb{C}} H^2(C, \mathbf{E}_{\rho_j}) = \chi(\pi_1(C)) = 2 - 2g .$$

Note that $H^0(C, \mathbf{E}_{\rho_j}) = 0$ because ρ_j is non-trivial. Furthermore, $H^1(C, \mathbf{E}_{\rho_j})$ is in a one-to-one correspondence with the set of non-equivalent representations $\rho : \pi_1(C) \longrightarrow T_2 \subset T_2(\mathbb{C})$ with the upper diagonal character ρ_j . Using the identity $\prod_{i=1}^g [e_i, e_{g+i}] = e$ for generators $e_1, \dots, e_{2g} \in \pi_1(C)$ we easily obtain that $\dim_{\mathbb{C}} H^1(C, \mathbf{E}_{\rho_j}) = 2g - 2$. Thus we have $H^2(C, \mathbf{E}_{\rho_j}) = 0$. This shows that

$H^2(\pi_1(C), \mathbb{C}) = 0$ and R_j is isomorphic to the semidirect product of $\mathbb{C}$ and $\pi_1(C)$, i.e., $R_j = \mathbb{C} \ltimes \pi_1(C)$ with multiplication

$$(v_1, g_1) \cdot (v_2, g_2) = (v_1 + \rho_j(g_1) \cdot v_2, g_1 \cdot g_2), \quad v_1, v_2 \in \mathbb{C}, \quad g_1, g_2 \in \pi_1(C).$$

Let us determine a map ϕ_j of R_j to T_2 by the formula

$$\phi_j(v, g) = \begin{pmatrix} \rho_j(g) & v \\ 0 & 1 \end{pmatrix}$$

Obviously, ϕ_j is a correctly defined homomorphism with upper diagonal character ρ_j . Hence $\phi_j \circ t_j : \pi_1(M) \rightarrow T_2$ is a homomorphism which is non-trivial on $\text{Ker}(f_*)$ by its definition. Now Proposition 2.3 implies that ρ_j is a unitary character. Therefore we have proved that all characters of the action of $\pi_1(C)$ on H are unitary. This means that each element of the above action is defined by a matrix from $SL_k(\mathbb{Z})$ with unitary eigen values. Applying to these eigen values the Kronecker theorem asserting that an algebraic integer is a root of unity if and only if all its Galois conjugates are unitary (see, e.g. [BS, p.105, Th.2]) we obtain that each ρ_j is a torsion character. $\square$

2.5. In this section we will prove a result about the structure of $\text{Ker}(f_*)$ where $f : M \rightarrow C$ is a surjective holomorphic map with connected fibres of a compact Kähler manifold M onto a curve C .

Using the Lojasiewicz triangulation theorem [L, Th.2], compactness of M and the fact that all fibres of f are connected, we can find an open finite cover $(U_i)_{1 \leq i \leq s}$ of C such that $W_i := f^{-1}(U_i)$ is an open neighbourhood of a fibre $V_i = f^{-1}(x_i)$, $x_i \in U_i$, and V_i is a deformation retract of some open $\widetilde{W}_i \supset W_i$. Let us fix some points $x_i \in V_i$ and paths γ_i connecting x_1 with x_i , $1 \leq i \leq s$. By G_i we denote the image of $\gamma_i \cdot \pi_1(V_i, x_i) \cdot \gamma_i^{-1}$ in $\pi_1(M, x_1)$. (Clearly, each G_i is finitely generated.)

Lemma 2.6 *$\text{Ker}(f_*)$ is the minimal normal subgroup of $\pi_1(M, x_1)$ containing all G_i ($1 \leq i \leq s$).*

Proof. Let $R \subset \pi_1(M, x_1)$ be the minimal normal subgroup containing all G_i . Clearly, $R \subset \text{Ker}(f_*)$. We will prove the reverse inclusion. Let $p : M_1 \rightarrow M$ be the regular covering over M with the transformation group $R_1 := \pi_1(M)/R$. It is a principle bundle over M with discrete fibre R_1 . We will prove that the restriction of M_1 to each $\widetilde{W}_i$ is the trivial bundle. Since V_i is a deformation retract of $\widetilde{W}_i$, it suffices to prove that for any closed path $\gamma \subset V_i$ passing through x_i and any point $y \in p^{-1}(x_i)$ the unique covering path $\gamma' \subset M_1$ of γ passing through y is closed. To prove this let us consider the path $s := \gamma_i \cdot \gamma \cdot \gamma_i^{-1}$. Let $s_1 \subset M_1$ be any path which covers s . Then s_1 is closed, because s represents an element of the *normal subgroup* $R \subset \pi_1(M)$. Let γ' be the covering of γ with endpoints y and y' . Then there is an element $g \in R_1$ such that $y' = g(y)$. Let h and h' be the unique covering paths of γ_i and γ_i^{-1} passing through y and y' , respectively. Clearly $h' = g(h^{-1})$. Now $s' = h \cdot \gamma' \cdot h'$ covers s and so it is closed. Then we have $h' = h^{-1}$ implying $g = 1$ and $y = y'$. So we proved that M_1 is trivial over each $\widetilde{W}_i$ and, in particular, it is trivial over each W_i . This means that M_1 is defined by a locally constant cocycle $\{c_{ij}\}$ (with

values in R_1) defined on the cover $(W_i)_{i \in I}$ of M . But then $\{c_{ij}\}$ is the pullback of a cocycle defined on $(U_i)_{i \in I}$ because the fibres of f are connected. This cocycle determines a principle bundle C_1 on C with the fibre R_1 such that $f^*C_1 = M_1$. In particular, C_1 determines a homomorphism $q : \pi_1(C) \rightarrow R_1$ such that $q \circ f_*$ is the quotient homomorphism $\pi_1(M) \rightarrow R_1$. This implies that $\text{Ker}(f_*) \subset R$. $\square$

2.6. We use Lemma 2.6 to prove the following result.

Lemma 2.7 *Let $\rho \in \text{Hom}(\pi_1(M), T_2)$ be such that $\text{Ker}(f_*) \subset \text{Ker}(\rho_a)$ where ρ_a is unitary. Then $\rho(\text{Ker}(f_*)) \subset N_2$ is a free abelian group of finite rank.*

Proof. Without loss of generality we will assume that $\text{Ker}(f_*) \not\subset \text{Ker}(\rho)$. Let $i : V \hookrightarrow \pi_1(M)$ be the fibre over a regular value of f and $H = i_*(\pi_1(V)) \subset \pi_1(M)$ be the corresponding normal subgroup. First, we prove that $\rho|_H$ is not trivial. Indeed, let E be the flat vector bundle associated to ρ_a . By our assumptions $E = f^*L$ where L is a flat vector bundle on C with unitary structure group. It is well known that the equivalence class of ρ is defined by a harmonic 1-form η with values in E . Since the structure group of E is unitary, the Hodge decomposition implies that $\eta = \omega_1 + \omega_2$, where ω_1 is holomorphic and ω_2 is antiholomorphic. From the proof of Lemma 2.4 we know that ρ_H is defined by integration of η along closed paths of $i(V)$. Assume to the contrary that $\rho|_H$ is trivial. This means that $\eta|_{i(V)}$ represents 0 in $H^1(i(V), \mathbb{C})$ and so $\omega_1|_{i(V)} = \omega_2|_{i(V)} = 0$. But then Lemma 2.2 implies that $\omega_i|_V = 0$ ($i = 1, 2$) for any fibre V of f . Let us consider a path γ which represents an element of $\pi_1(V_i, x_i)$ (the notation is as in Lemma 2.6). Let $\gamma' = \gamma_i \cdot \gamma \cdot \gamma_i^{-1}$. Then the monodromy argument shows that $\rho(\gamma') = C \cdot A \cdot C^{-1}$ where $C \in T_2$, $A \in N_2$ and A is defined by integration of η along γ . Therefore $\rho(\gamma') = 1 \in N_2$. Let $[\gamma'] \in \pi_1(M)$ be the element defined by γ' . According to Lemma 2.6, $\text{Ker}(f_*)$ is the minimal normal subgroup containing all such $[\gamma']$ (for any i). The above argument then shows that $\rho|_{\text{Ker}(f_*)}$ is trivial which contradicts to our assumption.

Next, we will prove that $\rho_a(\pi_1(M))$ consists of algebraic integers. Let $H_f \cong \mathbb{Z}^k$ be the free part of $H^{ab} = H/DH$. In what follows we identify H_f with $\mathbb{Z}^k$. Let $s : \pi_1(M) \rightarrow SL_k(\mathbb{Z})$ be the representation induced by the action of $\pi_1(M)$ on H by conjugation. Since $\rho(H_f) = \rho(H)$ is non-trivial and consists of unipotent matrices, $k \geq 1$ and identity (2.1) implies that the action of $\rho(\pi_1(M))$ on $\rho(\mathbb{Z}^k)$ by conjugation is defined by multiplication of non-diagonal elements of $\rho(\mathbb{Z}^k)$ by elements of $\rho_a(\pi_1(M))$. Let $e_1, \dots, e_k$ be the standard basis in $\mathbb{Z}^k$. Identifying N_2 with $\mathbb{C}$ we obtain a nonzero vector $v = (\rho(e_1), \dots, \rho(e_k)) \in \mathbb{C}^k$. Then a simple calculation shows that v is a nonzero eigen vector of $s(g)$ with the eigen value $\rho_a(g)$, $g \in \pi_1(M)$. This completes the proof.

Now let us finish the proof of the lemma. Let $\{g_j\}$, $1 \leq j \leq l$, be the *finite family* which consists of all generators of all G_i (as in Lemma 2.6). Then according to Lemma 2.6, $\rho(\text{Ker}(f_*)) \subset \rho(\pi_1(M))$ is the minimal normal subgroup containing all $h_j := \rho(g_j)$. Since N_2 is abelian, identity (2.1) shows that $\rho(\text{Ker}(f_*))$ is a discrete abelian group generated by all elements $\rho_a(g) \cdot g_j$, $g \in H_1(M, \mathbb{Z})$, $1 \leq j \leq l$. Let $t_1, \dots, t_p$ be the family of generators of $H_1(M, \mathbb{Z})$. Then the above arguments imply that each $\rho_a(t_i), \rho_a(-t_i)$ ($i = 1, \dots, p$) are roots of polynomials of degree k with integer coefficients and leading coefficients 1. Since any $g \in H_1(M, \mathbb{Z})$ can be

written as $\sum_{i=1}^p a_i t_i + \sum_{i=1}^p b_i (-t_i)$ with $a_i, b_i \in \mathbb{Z}_+$, the previous shows that the abelian group $\rho(Ker(f_*))$ is generated (over $\mathbb{Z}$) by the elements

$$\rho_a(t_1)^{a_1} \dots \rho_a(t_p)^{a_p} \cdot \rho_a(-t_1)^{b_1} \dots \rho_a(-t_p)^{b_p} \cdot g_j$$

where a_i, b_i are nonnegative integers satisfying $0 \leq a_i \leq k-1$, $0 \leq b_i \leq k-1$, and $1 \leq j \leq l$. Since the number of such generators is finite, $\rho(Ker(f_*))$ is finitely generated. It is also free as a subgroup of the free group N_2 .

The proof of the lemma is complete. $\square$

2.7. Proof of Theorem 2.1. Let $\rho \in Hom(\pi_1(M), T_2)$ be such that $Ker(f_*) \subset Ker(\rho_a)$ but $Ker(f_*) \not\subset Ker(\rho)$. According to Proposition 2.3, ρ_a is unitary. Then according to Lemma 2.7, $\rho(Ker(f_*)) \subset N_2$ is isomorphic to $\mathbb{Z}^r$, $r \geq 1$. We set $G = Ker(f_*) \cap Ker(\rho)$ and $R := \pi_1(M)/G$. Then R is defined by the sequence

$$\{1\} \longrightarrow Ker(f_*)/G \longrightarrow R \longrightarrow \pi_1(C) \longrightarrow \{1\}.$$

where $Ker(f_*)/G \cong \rho(Ker(f_*)) = \mathbb{Z}^r$. Since $\rho(Ker(f_*)) \subset N_2$, the above definition implies that the natural action of $D\pi_1(C)$ on $Ker(f_*)/G$ is trivial. Thus according to Proposition 2.5, all characters of the induced representation $\pi_1(C) \rightarrow GL_r(\mathbb{C})$ are torsion. In particular, ρ_a is torsion as one of such characters.

The proof of the theorem is complete. $\square$

References

- [A] M.F.Atiyah, Complex analytic connections in fibre bundles, Trans. Amer. Math. Soc. **85** (1957), 181-207.
- [Be] A.Beauville, Annulation du H^1 pour les fibrés en droites plats, Lecture Notes in Math., **1507**, Springer Verlag (1992), 1-15.
- [BS] A.I.Borevich and I.R.Shafarevich, Number theory, Translated from Russian. Pure and Applied Mathematics, vol. 20, Academic Press, New York, 1966.
- [Br] A.Brudnyi, Classification theorem for a class of flat connections and representations of Kähler groups, Michigan Math. J., **46** (3) (1999), 489-514.
- [Ca] F.Campana, Ensembles de Green-Lazarsfeld et quotients résolubles des groupes de Kähler, J. Alg. Geom. **10** (2001), no. 4, 599-622.
- [G] A. Guichardet, Cohomologie des groupes topologiques et des algèbres de Lie, Cedec/Fernand Nathan, Paris (1980).
- [L] S. Lojasiewicz, Triangulation of semi-analytic sets, Ann. Scuola Norm. Sup. Pisa, (3) **18** (1964), 449-474.
- [M] D. Mumford, Abelian varieties, Lect. in Tata Inst. of Fundamental Research, Bombay (1968).
- [S] C.T.Simpson, Subspaces of Moduli Spaces of rank one local systems, Ann. Sc. ENS, **26** (1993), 361-401.